

AD_____

Award Number: DAMD17-96-1-6330

TITLE: Sex Differences in Susceptibility and Resistance to
Noise-Induced Hearing Loss in Chinchillas

PRINCIPAL INVESTIGATOR: Sandra L. McFadden, Ph.D.

CONTRACTING ORGANIZATION: State University of New York at Buffalo
Amherst, New York 14228-2567

REPORT DATE: October 1999

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20000828 201

DTIC QUALITY INSPECTED 4

REPORT DOCUMENTATION PAGEForm Approved
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE October 1999	3. REPORT TYPE AND DATES COVERED Annual (23 Sep 98 - 22 Sep 99)	
4. TITLE AND SUBTITLE Sex Differences in Susceptibility and Resistance to Noise-Induced Hearing Loss in Chinchillas			5. FUNDING NUMBERS DAMD17-96-1-6330	
6. AUTHOR(S) Sandra L. McFadden, Ph.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) State University of New York at Buffalo Amherst, New York 14228-2567 E-MAIL: mcfadden@acsu.buffalo.edu			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; Distribution Unlimited				12b. DISTRIBUTION CODE
13. ABSTRACT (Maximum 200 Words) There is tremendous variability in the amount of hearing loss individuals develop from a given noise exposure. The reasons for individual differences in susceptibility to noise-induced hearing loss (NIHL) are largely unknown, but may include factors such as levels of endogenous antioxidant enzymes or steroid hormones. Our experiments are focusing on the effects of steroid hormones on susceptibility to NIHL in chinchillas. Chinchillas treated with 17- β -estradiol prior to noise exposure developed less permanent hearing loss than controls. The protective effects of estradiol were found in two separate experiments, one utilizing a continuous noise (octave band noise with a center frequency of 4 kHz, at 105 dB SPL for 4 hours), and the other an impulse noise simulating M16 rifle fire. These experiments support the hypothesis that endogenous levels of estrogen could influence individual susceptibility to NIHL.				
14. SUBJECT TERMS Acoustic trauma; Gender differences; Hearing impairment; Noise-induced hearing loss; Steroid hormones; Estrogen; Estradiol				15. NUMBER OF PAGES 53
				16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

N/A Where copyrighted material is quoted, permission has been obtained to use such material.

N/A Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

SM Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

SM In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NIH Publication No. 86-23, Revised 1985).

N/A For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

N/A In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

N/A In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

N/A In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

Sandra L. McGladde 9-28-99
PI - Signature Date

TABLE OF CONTENTS

Front Cover	1
Standard Form 298, Report Documentation Page	2
Foreword	3
Table of Contents	4
1. INTRODUCTION	5
2. BODY	5-13
3. KEY RESEARCH ACCOMPLISHMENTS	13-14
4. REPORTABLE OUTCOMES	14
5. CONCLUSIONS	14
6. REFERENCES	15
7. APPENDICES	16

1. INTRODUCTION

All tasks of Technical Objective #1 in the approved Statement of Work were completed during the first two years of the project. In the third and final year of the project, significant progress was made on experiments outlined under Technical Objective #2. A one-year no-cost extension of the grant has been received, so that all tasks of Technical Objective #2 can be completed as outlined in the Statement of Work. The experiments currently being conducted are designed to explore the effects of steroid hormones on noise-induced threshold shifts and hair cell losses in chinchillas. This report describes the advances made toward fulfilling the tasks of Technical Objective #2 for the funding period from Sept. 23, 1998 to Sept. 22, 1999.

Specifically, this report describes experiments in which chinchillas were given estrogen (17- β -estradiol) prior to noise exposure. The results show that estrogen treatment reduces threshold shifts associated with exposure to two different types of noise, impulse noise simulating M16 rifle fire and continuous noise. The findings from the chinchilla offer insights into factors that may contribute to individual differences in noise susceptibility in humans.

2. BODY

Summary of task objectives for Technical Objective #2: Test auditory sensitivity before and after noise exposure in animals treated with steroid hormones, and examine the effects of steroid hormones on noise-induced hair cell loss.

2.1. METHODS

All procedures were reviewed and approved by the University of Buffalo Animal Care and Use Committee, and conformed to NIH guidelines for the humane treatment of laboratory animals.

2.1.1. Subjects and surgery

Subjects were 37 chinchillas (*Chinchilla langier*) between 1 and 3 years of age. Animals were deeply anesthetized with an intramuscular injection of ketamine hydrochloride (Ketaset; 56 mg/kg) and acepromazine (Promace; 0.56 mg/kg). Tungsten electrodes were implanted into the right and/or left inferior colliculus (IC) and the rostral cranium for recording auditory evoked potentials (EVPs). Details of the surgical procedures are provided in Appendix I (McFadden et al., 1999). Following surgery, the animals recovered in a quiet animal colony for at least one week prior to testing.

2.1.2. EVP testing

The auditory sensitivity of each animal was determined from EVP thresholds. All testing was conducted in a sound attenuating booth (Industrial Acoustics Corp. 400) lined with sound absorbing foam panels. The awake chinchilla was placed in a custom-designed tube (Snyder and Salvi, 1994) that held its head at a constant orientation within the calibrated sound field. Stimuli consisted of 10 ms tones (2 ms cosine rise/fall ramp, alternating phase) at octave intervals from 0.5 to 16 kHz, presented at a rate of 19/sec. Stimulus level was incremented in 5 dB steps from below threshold to 80 dB SPL. Details of the stimuli and recording procedures can be found in Appendix I (McFadden et al., 1999).

2.1.3. Noise stimuli and acoustic calibration

Animals were exposed to either impulse noise or octave band noise (OBN) with a center frequency of 4 kHz. The impulse noise simulated impulses created by a U.S. Army M16A1 rifle (Danielson, Henderson, Gratton, Bianchi, and Salvi, 1991). The impulses were generated digitally, attenuated (HP 350D), amplified (NAD 2200), and delivered to a compression driver (JBL 2446) coupled to a sound delivery tube (5 cm dia X 20 cm) whose end was cut at a 45° angle to broaden the range of resonance (Danielson et al., 1991). An animal in a restraint tube was placed 10 cm away from the acoustic driver and exposed to 50 pairs of impulses (100 total). Impulses in each pair were spaced 50 ms apart, and there was a 1000 ms interval between the onset of each pair (Henselman et al., 1994). The duration of the exposure was 50 sec. For calibration of the impulse noise, a 1/8" microphone (Bruel and Kjaer Model 4138) was placed at the position that would be occupied by a restrained animal. The voltage corresponding to a 114 dB, 250 Hz tone produced by a pistonphone coupled to the microphone was determined, and used to calculate the desired voltage for a 150 dB peak SPL signal. The attenuation was adjusted to produce the desired voltage.

The 4 kHz OBN was generated digitally, attenuated (HP 350D), amplified (NAD 2200), and delivered to a compression driver (JBL 2446) suspended from the ceiling of the sound booth. During exposure, animals were placed in individual cages beneath the loudspeaker and provided free access to food and water. The exposure level, calibrated with a Type I sound level meter and condenser microphone, was 105 dB SPL. Exposure duration was 4 hr.

2.1.4. Experimental protocol

Animals were randomly assigned to estradiol treatment groups, a vehicle control group, or an untreated control group. Animals in estradiol treatment groups received daily subcutaneous injections of 17- β -estradiol (Sigma Chemicals) dissolved in olive oil vehicle for 7-14 consecutive days. Animals in the vehicle control group received an equal volume of vehicle alone on the same schedule as estradiol-treated animals. IC-EVP thresholds were measured twice before treatment, and at various times relative to treatment and noise exposure. Post-noise measurements were made at 15 min, 24 hr, 7 days, and 14-21 days.

After all measurements had been obtained, chinchillas were deeply anesthetized with sodium pentobarbital (Somlethal, 100 mg/kg i.p.) and decapitated. The cochleas were quickly removed, stained with a succinate dehydrogenase (SDH) staining solution, and post-fixed with 10% formalin (see details in Appendix I, McFadden et al., 1999). Cochleas were dissected from the apex to the base, mounted in sections in glycerin on microscope slides, coverslipped, and examined using light microscopy (400X magnification). The numbers of missing OHCs and IHCs were determined for successive segments of the organ of Corti. Individual cochleograms were constructed to show the percentage of hair cells missing as a function of distance from the apex of the cochlea. Percent hair cells missing was referenced to our lab standards based on average hair cell counts from 9 cochleas of young (<1 yr old), healthy chinchillas. Percent distance from the apex was converted to frequency using the frequency-place map of Greenwood (1990).

2.1.5. Data analyses

Data analyses were geared toward answering two questions. First, does estradiol treatment affect basic auditory sensitivity? Second, does estradiol influence susceptibility to impulse noise or continuous noise? Analyses of variance (ANOVAs) and Student t-tests were used to assess differences between means. The dependent variables were IC-EVP thresholds and threshold shifts at various times after treatment or noise exposure. All statistical tests were evaluated using a 0.05 criterion of significance.

2.1.6. Estradiol assays

Blood samples were collected from deeply anesthetized chinchillas prior to cochlear histology. The blood samples were centrifuged to separate serum for estradiol assays. Samples were treated with a steroid displacement reagent to free estradiol bound to transport proteins in the serum. Estradiol levels were measured using an Enzyme Immunoassay kit (Assay Designs Inc., product number 90108). Microplates from the assay were read using a microplate reader from Bio-Rad Laboratories, Inc. (Model 3550-UV). All samples were run in triplicate to ensure reliability.

2.2. RESULTS AND DISCUSSION**2.2.1. Preliminary results of estradiol assays**

The following table presents measured levels of estradiol in picograms/ml serum. For each subject listed, the value is the average of 3 assays.

Females

109.01
256.58
258.36
263.92
312.76

M=240.12; sd= 76.87
Males

40.82
66.92
69.98
82.40

M=65.03; sd= 17.47


Two aspects of these results are particularly interesting. First, the levels for female and male chinchillas are in the range reported for humans. This supports the use of the chinchilla in studies of hormone effects on hearing. Second, females have higher levels and are much more variable than males. Because individuals show a wide range of variability, it will be possible in

future studies to correlate endogenous levels of hormones with susceptibility to NIHL in both treated and untreated populations.

2.2.2. Experiments with estradiol pre-treatment


The effects of 17- β -estradiol on susceptibility to NIHL were studied in two experiments. In Experiment I, chinchillas in the estradiol groups were given daily injections of estradiol in olive oil for 1-2 weeks before exposure, for total doses of 200-265 mg. Animals in a vehicle control group received injections of the olive oil vehicle alone. Thresholds measured on Days 2, 4 and 7 during the course of hormone treatment were not different from those measured before treatment, indicating that short-term estradiol treatment has no direct effect on auditory sensitivity. All animals were exposed to impulse noise, and EVPs were measured 15 min, 24 hr, 7 days and 14 days after exposure.

Permanent threshold shifts are shown in Figure 1. The animals in the vehicle control group had PTS ranging from 20 dB at 0.5 kHz to 55 dB at 8 kHz. Chinchillas pre-treated with estradiol had significantly less PTS at frequencies between 2 and 8 kHz. On average, chinchillas pre-treated with estradiol had PTS values ranging from 15 dB at 0.5 kHz to 30 dB at 2 kHz. At 8 kHz, estradiol-treated chinchillas had less than 20 dB hearing loss—a “savings” of 35 dB relative



to controls. These results indicate that estradiol treatment provides protection from threshold shifts caused by impulse noise.

Hair cell losses are shown in Figure 2. Inner hair cell losses are shown in the top panel; outer hair cell losses are compared in the bottom panel. Overall, there was no dramatic difference in hair cell loss between estradiol-treated animals and the vehicle control animals. Although




estradiol-treated animals had much less hair cell loss in the basal-most 10-20% of the cochlea, differences in other regions were minor. This finding is interesting because it suggests that estradiol may act on the stria vascularis rather than the organ of Corti. However, it should be noted that it is not unusual to see a lack of correspondence between hair cell loss and hearing loss

(Hamernik et al., 1989; Boettcher et al., 1992; McFadden et al., 1997), and the meaning of this dissociation is not clear.

In Experiment II, chinchillas in the estradiol groups received estradiol for 7 days prior to exposure. Total estradiol dose was either 100 μg (N=5) or 725 μg (N=6). Animals in a control group received no treatment prior to noise. All animals were exposed to 4 kHz octave band noise at a level of 105 dB SPL for 4 hours. This is the same exposure that was used in a recent study of protection using the chemical R-phenylisopropyladenosine (R-PIA) (Hu et al., 1997). In that study, R-PIA was placed on the round window of the cochlea and wicked off after 30 minutes. Chinchillas treated with R-PIA developed significantly less PTS than controls that had received a drop of saline on the round window. The results from the Hu et al. experiment are compared to the current results to provide a perspective on the magnitude of protection afforded by systemic treatment with estradiol.

Figure 3 shows recovery of hearing over time for chinchillas treated with estradiol. Threshold shifts at 14 days represent PTS. The recovery curves are very similar for animals in the low dose



and the high dose groups. However, animals in the high dose group had less threshold shift at high frequencies than animals in the low dose group. The greatest divergence between treatment groups is seen at 4 and 8 kHz on Day 7. Differences at 4 kHz persist at Day 14. Figure 4 shows PTS as a function of frequency for treated animals and controls. Results from Hu et al. (1997) are shown for comparison. It is clear that estradiol treatment reduced PTS relative to controls. The high dose of estradiol produced savings equivalent to those seen following R-PIA treatment in the Hu et al. (1997) experiment. This is important, because the protective effects achieved with R-PIA involved invasive surgery, whereas the protective effects of estradiol were achieved with simple systemic treatment.




Figure 4. Permanent threshold shifts caused by 4 kHz octave band noise.

Hair cell losses are shown in Figure 5. As with the previous experiment (Fig. 2), there were no remarkable differences between treated and control groups in inner hair cell loss (upper panel) or outer hair cell loss (lower panel). These results are consistent with the hypothesis that estradiol acts on vascular tissue of the cochlea rather than on the hair cells themselves.




Figure 5. Hair cell losses after exposure to 4 kHz octave band noise.

Like R-PIA, steroid hormones can act through many different routes, each of which could be important for modulating susceptibility to NIHL. Estradiol can potentiate the activity of gamma-aminobutyric acid, a ubiquitous inhibitory neurotransmitter of the central nervous system; it can affect neuronal activity via changes in cellular neurochemistry and morphology; it can act on cell membranes to alter permeability to neurotransmitters, precursors and receptor functioning; it can act directly as an antioxidant; and it can influence the bioactivity of other antioxidants and blood flow promoters such as nitric oxide (Arnal et al., 1996; Ayres et al., 1996; Behl et al., 1995; Chadwick and Widdows, 1990; Goodman et al., 1996; Romer et al., 1997; Ruiz-Larrea et al., 1994).

2.2.3. PROBLEMS

Progress toward completing all experiments outlined for the third year of the grant was impeded by several unforeseeable problems. Most of these revolved around animal availability and health. The first animal problem arose when our regular chinchilla supplier ran out of chinchillas early in the year, forcing us to locate a different supplier. The first batch of animals we purchased from the new breeder was of excellent quality, but subsequent batches have been of very mixed quality, and a large number of animals have been unsuitable for our experiments. Our regular supplier now has a minimum order size of 30. This limits how frequently we can order animals, due to our limited cage space. We have also had problems getting an adequate number of female chinchillas for our experiments.

Progress was also slowed down by changes in personnel and the need to develop new techniques for assaying hormone levels from serum samples. During the course of the year, two new surgeons were trained to perform the electrode implantation surgeries, a new technician was trained for histological evaluations of cochleas, and two technicians were trained in animal handling and testing procedures.

Due to the problems described above, we applied for and received a one-year no-cost extension on the grant. This additional year will allow us to finish all experiments as originally outlined in our Statement of Work.

3. KEY RESEARCH ACCOMPLISHMENTS—Year 3

- * Estradiol levels were determined from serum samples of female and male chinchillas, and found to vary on a continuum comparable to humans.
- * The protective effects of estradiol on noise-induced hearing loss were demonstrated in two separate experiments that utilized different noise exposure conditions. Estradiol reduced NIHL from octave band noise centered at 4 kHz and impulse noise simulating M16 rifle fire.
- * Estradiol was determined to have protective effects similar to another known otoprotectant, R-PIA. Unlike R-PIA, however, the protective effects of estradiol were achieved with simple systemic treatment rather than invasive surgery.

* The mechanism of estradiol protection does not appear to involve sparing of cochlear hair cells. Estradiol may exert its effects on the stria vascularis, or by acting as an antioxidant or a modulator of cochlear neurotransmitter substances.

4. REPORTABLE OUTCOMES: Manuscripts, abstracts & presentations

1. McFadden, S.L., Henselman, L.W., and Zheng, X.Y. (1999). Sex differences in auditory sensitivity of chinchillas before and after exposure to impulse noise. *Ear Hear.* **20**, 164-174.
2. McFadden, S.L., Zheng, X.Y., and Ding, D.L. Conditioning-induced protection from impulse noise in female and male chinchillas. *J. Acoust. Soc. Am.* (submitted 7/26/99).
3. McFadden, S.L., Zheng, X.Y., and Ding, D.L. Sex differences in threshold shifts and hair cell loss in chinchillas exposed to simulated military noises. (manuscript in preparation; to be submitted to *Noise and Health*)
4. McFadden, S.L. Overview of research on steroid hormones and noise-induced hearing loss. *Lake Ontario Hearing Meeting*, Syracuse University, June 15, 1999.
5. McFadden, S.L., Zheng, X.Y., Ding, D.L., and Henderson, D. (1999) Differences between female and male chinchillas in susceptibility to noise-induced hearing loss. *Assoc. Res. Otolaryngol. Abstr.* **610**.
6. Lockwood, D., McFadden, S.L., Jiang, H., and Rosenberg, L. Systemic treatment with estradiol reduces noise-induced hearing loss in the chinchilla. *Assoc. Res. Otolaryngol. Abstr.* (submitted 10/1/99).
7. McFadden, S.L. and Henderson, D. (1999) Recent advances in understanding and preventing noise-induced hearing loss. *Current Opin. Otolaryngol. Head Neck Surg.*, **7**(5).

5. CONCLUSIONS

The results from the experiments described here have important practical and theoretical implications. From a theoretical standpoint, they provide a much-needed perspective on the role of steroid hormones on normal physiology and function, thereby increasing our understanding of basic auditory system physiology. Gender differences have frequently been reported in humans, but the basis for these differences has remained elusive. Often, sex differences have been attributed to differences in noise exposure history, as males are more often engaged in occupational and recreational activities that involve high-level noise. However, our findings argue for more fundamental, inherent differences related to biochemical and hormonal factors. Based on our findings, it is reasonable to hypothesize that individuals with high estrogen levels will be less susceptible to NIHL than individuals with low estrogen levels.

6. REFERENCES

- Arnal, J.F., Clamens, S., Pechet, C., Negre-Salvayre, A., Allera, C., Girolami, J.-P., Salvayre, R. and Bayard, F. (1996) Ethinylestradiol does not enhance the expression of nitric oxide synthase in bovine endothelial cells but increases the release of bioactive nitric oxide by inhibiting superoxide anion production. *Proc. Natl. Acad. Sci. USA* 93, 4108-4113.
- Ayres, S., Tang, M. and Subbiah, M.T.R. (1996) Estradiol-17 β as an antioxidant: Some distinct features when compared with common fat-soluble antioxidants. *J Lab Clin Med* 128, 367-375.
- Behl, C., Widmann, M., Trapp, T. and Holsboer, F. (1995) 17- β estradiol protects neurons from oxidative stress-induced cell death *in vitro*. *Biochem Biophys Res Comm* 216, 473-482.
- Boettcher, F.A., Spongr, V.P. and Salvi, R.J. (1992) Physiological and histological changes associated with the reduction in threshold shift during interrupted noise exposure. *Hear Res* 62, 217-236.
- Danielson, R., Henderson, D., Gratton, M. A., Bianchi, L. and Salvi, R. (1991) The importance of "temporal pattern" in traumatic impulse noise exposures. *J Acoust Soc Am* 90, 209-218.
- Goodman, Y., Bruce, A.J., Cheng, B. and Mattson, M.P. (1996) Estrogens attenuate and corticosterone exacerbates excitotoxicity, oxidative injury, and amyloid β -peptide toxicity in hippocampal neurons. *J Neurochem* 66, 1836-1844.
- Greenwood, D.D. (1990) A cochlear frequency-position function for several species—29 years later. *J Acoust Soc Am* 87, 2592-2604.
- Hamernik, R.P., Patterson, J.H., Turrentine, G.A., and Ahroon, W.A. (1989) The quantitative relation between sensory cell loss and hearing thresholds. *Hear Res* 38, 199-212.
- Henselman, L.W., Henderson, D., Subramaniam, M. and Sallustio, V. (1994) The effect of 'conditioning' exposures on hearing loss from impulse noise. *Hear Res* 78, 1-10.
- Hu, B.H., Zheng, X.Y., McFadden, S.L., Kopke, R. and Henderson, D. (1997) R-PIA attenuates noise-induced hearing loss in the chinchilla. *Hear Res* 113, 198-206.
- Romer, W., Oettel, M., Driescher, P. and Schwarz, S. (1997) Novel "scavestrogens" and their radical scavenging effects, iron-chelating, and total antioxidative activities: delta-8,9-dehydro derivatives of 17 α -estradiol and 17 β -estradiol. *Steroids* 62, 304-310.
- Ruiz-Larrea, M.G., Leal, A.M., Liza, M., Lacort, M. and de Groot, H. (1994) Antioxidant effects of estradiol and 2-hydroxyestradiol on iron-induced lipid peroxidation of rat liver microsomes. *Steroids* 59, 383-511.
- Snyder, D.L. and Salvi, R.J. (1994) A novel chinchilla restraint device. *Lab Anim* 23, 42-44.

LIST OF APPENDICES

- I. Paper published in *Ear and Hearing*.
- II. Abstract submitted to Association for Research in Otolaryngology, 2000.
- III. Poster presented at Association for Research in Otolaryngology, 1999.
- IV. Manuscript submitted to *Journal of the Acoustical Society of America*.

Sex Differences in Auditory Sensitivity of Chinchillas Before and After Exposure to Impulse Noise

Sandra L. McFadden, Lynn W. Henselman, and Xiang-Y. Zheng

Objective: To determine if chinchillas exhibit sex differences in 1) basic auditory sensitivity and 2) susceptibility to cochlear damage and hearing loss from high-level impulse noise.

Design: The auditory sensitivity of 73 chinchillas was assessed by measuring evoked potentials from electrodes implanted in the inferior colliculus (IC-EVPs) and cubic ($2f_1$ - f_2) distortion product otoacoustic emissions (CDPs). A subgroup of 16 chinchillas were retested after exposure to simulated M16 rifle fire (150 dB pSPL impulse noise). Thresholds and postexposure temporary and permanent threshold shifts were compared as a function of sex and frequency using analysis of variance procedures. Cochleograms, showing the percent of hair cells missing as a function of location on the basilar membrane, were constructed to show inner hair cell (IHC) and outer hair cell (OHC) losses for each group.

Results: Female chinchillas had slightly lower high-frequency thresholds, and slightly higher low-frequency thresholds than male chinchillas, but similar IC-EVP and CDP amplitude functions. Significant sex differences were observed after exposure to high-level impulse noise. Overall, female chinchillas developed approximately 10 dB more high-frequency hearing loss, but approximately 5 dB less low-frequency hearing loss than males. Hair cell losses, particularly IHC losses, were substantially less for females as compared with males.

Conclusions: The results point to close similarities between chinchillas and humans with regard to sex/gender differences in basic auditory sensitivity before noise exposure, suggesting that the chinchilla may be a good model for exploring the anatomical and physiological bases of these differences. In addition, the results show significant sex differences in the physiological and anatomical response of the chinchilla cochlea to high-level noise. Similar differences in humans could have important implications with regard to military assignments and hearing conservation programs.

(*Ear & Hearing* 1999;20;164-174)

Center for Hearing and Deafness (S.L.M., X.Y.Z.), University of Buffalo, Buffalo, New York; and Office of the Inspector General (L.W.H.), Department of Defense, Arlington, Virginia.

Noise-induced hearing loss (NIHL) is a major occupational hazard for military personnel because of the types and levels of noise encountered in training and combat situations (Dancer & Franke, 1986; Henselman, Henderson, Shadoan, Subramaniam, Saunders, & Ohlin, 1995; Henselman, Henderson, Subramaniam, & Sallustio, 1994). Damage to the cochlea can be caused by a variety of acoustic events, ranging from prolonged exposure to continuous noises that cause metabolic and biochemical changes in the cochlea, to relatively brief exposures to high-level impact and impulse noises such as gunfire, cannon fire and explosions, that can produce direct mechanical damage as well (Dancer & Franke, 1986; Henderson, Hamernik, & Sitler, 1974; Henderson, Spongr, Subramaniam, & Campo, 1994). A recognition of the serious consequences of NIHL led the U.S. Air Force to develop the first hearing conservation program (HCP) in 1948. The U.S. Navy and U.S. Army developed similar HCPs in 1955 and 1956, respectively (Henselman et al., 1995). Since their inception, military HCPs have served to increase awareness of the damaging effects of high-level noise exposure. They have also served to reduce the incidence and magnitude of NIHL in military personnel, primarily by mandating the use of personal protection devices such as sound-attenuating ear plugs or earmuffs in high-noise situations. However, NIHL remains a serious problem for military personnel who are exposed to loud noises during training and combat situations in which personal protection devices are either unavailable, impractical or dangerous to use, improperly fitted or worn, or inadequately designed to protect the ear from damage (Dancer, Buck, Parmentier, & Hamery, 1998).

As women become more fully integrated into a variety of military occupational specialties, many will be placed at risk for developing NIHL. It is critical, therefore, that we understand the specific relationship between noise exposure and hearing loss in women, so that appropriate measures for preventing NIHL can be developed and implemented.

Previous studies (Chung, Mason, Gannon, & Willson, 1983; Corso, 1963; Pearson et al., 1995; Ward,

1966) have reported small differences (generally less than 3 dB) between males and females in auditory sensitivity (i.e., thresholds for detecting pure tones under quiet listening conditions). In general, females tend to have slightly better pure-tone thresholds than males at frequencies above 1 to 2 kHz, whereas males may have slightly better thresholds below 1 to 2 kHz. Small, but consistent gender differences* have also been reported in susceptibility to temporary threshold shifts (TTS) caused by exposure to continuous tones or noise (Axelsson & Lindgren, 1981; Dengerink, Dengerink, Swanson, Thompson, & Chermak, 1984; Petiot & Parrot, 1984; Ward, 1966). In general, experimental studies of TTS in humans have found that males exhibit more TTS than females from low-frequency exposures (below 2 kHz), whereas females exhibit more TTS than males from high-frequency exposures (above 2 kHz). In an early investigation of gender differences in susceptibility to TTS produced by high intensity tones and noise, Ward (1966) conducted 17 experiments with 24 male and 25 female adults. Females showed approximately 30% less TTS than males when the exposure frequency was below 1 kHz, but approximately 30% more TTS when the exposure frequency was above 2 kHz.

The above studies examined TTS rather than the more important issue of permanent threshold shift (PTS) because it is not ethical to intentionally induce PTS in human subjects. Most of what little we know about gender differences in PTS comes from retrospective studies of workers exposed to noise in industrial settings (Berger, Royster, & Thomas, 1978; Gallo & Glorig, 1964). Under these conditions, which typically involve exposure to low-frequency continuous noises, males tend to develop much more hearing loss than females. Both Berger et al. (1964) and Gallo and Glorig (1964) found approximately 20 dB more PTS in males than in females after 9 yr of industrial noise exposure. These results are consistent with the gender differences observed in Ward's (1966) studies of TTS. However, there are no comparable studies of gender differences in PTS produced by exposures to high-level impulse noises that are typically found in military environments. A finding of gender differences in susceptibility to NIHL could have important implications for military assignments and HCPs.

The present study was conducted to determine whether there are systematic differences between female and male chinchillas in 1) basic auditory function, as assessed by inferior colliculus evoked

potentials (IC-EVPs) and cubic ($2f_1-f_2$) distortion product otoacoustic emissions (CDPs), and 2) their susceptibility to high-level impulse noise. Basic auditory function was assessed in a relatively large group of chinchillas ($N = 73$). Susceptibility to impulse noise was examined in a subgroup of these animals ($N = 16$). Findings from the chinchilla may shed light on gender differences in susceptibility to impact/impulse noise, and offer insights into the anatomical and physiological mechanisms contributing to documented gender differences in humans.

METHODS

All procedures described here were reviewed and approved by the University of Buffalo Animal Care and Use Committee, and conformed to federal guidelines for the humane treatment of laboratory animals.

Subjects

A total of 73 chinchillas (*Chinchilla laniger*; 37 female, 36 male) between 1 and 3 yr of age were used. A subgroup of animals (eight female, eight male) was exposed to impulse noise, and their thresholds were measured at various times after exposure (see below). The chinchilla was used for these studies because it 1) is relatively immune to middle ear infections and diseases that affect hearing (Clark, 1984); 2) has a relatively long life span (12 to 20 yr) with minor age-related cochlear pathology and hearing loss before 8 to 10 yr of age (Bohne, Gruner, & Harding, 1990; McFadden, Campo, Quaranta, & Henderson, 1997); and 3) reacts predictably to anesthesia and tolerates surgery well. Most importantly, the chinchilla has a range of hearing that is more similar to that of humans than most other laboratory animals, particularly in the low frequencies (Heffner & Heffner, 1991; Miller, 1970), which enhances its suitability as a model for studying NIHL (McFadden, Campo, Ding, & Quaranta, 1998). With regard to size, Clark (1984) states that female chinchillas tend to be larger than males. In a small group of our chinchillas (eight female, eight male) for which reliable weights were available, weight differences were minor, but favored females. The average weight of females was 572.2 g ($SD = 73.7$), versus 563.9 g ($SD = 70.0$) for males.

Surgical Preparation

Each animal was deeply anesthetized with an intramuscular injection of ketamine hydrochloride (60 mg/kg) and acepromazine (0.5 mg/kg). Chronic recording electrodes were implanted in the left and/or right inferior colliculus (IC), and in the ros-

* The term "gender differences" will be used to refer to male/female differences in humans, and the term "sex differences" will be used to refer to male/female differences in chinchillas and other nonhuman species.

tral cranium (McFadden et al., 1997). Thirteen animals were implanted unilaterally; all others were implanted bilaterally. A small hole was drilled in the dorsal cranium overlying the IC, and a recording electrode mounted on a stereotaxic device was advanced through the IC while the surgeon monitored sound-evoked electrical activity on audio and video monitors. When the electrode had been advanced to a depth that produced clear, large amplitude EVPs, it was cemented to the skull with cyanoacrylic adhesive and dental cement. A short electrode was implanted in the rostral cranium to serve as the common lead for IC-EVP recording. Because the electrodes remain fixed in position, variability associated with changes in electrode placement between tests is eliminated. In addition, the signal to noise ratio is much better with implanted electrodes than with more conventional scalp electrodes, so that thresholds can be determined with greater precision. IC-EVPs recorded from electrodes implanted in this manner yield thresholds that are very close to behavioral thresholds measured in the same animals (Henderson, Hamernik, Salvi, & Ahroon, 1983), and about 15 to 20 dB lower than threshold estimates obtained using subcutaneous electrodes in the same animals (Murphy & Themann, Reference Note 1). After surgery, the animals recovered in a quiet animal colony for at least 1 wk before testing.

Measures of Auditory Function

The auditory sensitivity of each animal was assessed by measuring IC-EVPs. CDPs were also obtained from most animals. All testing was conducted in a sound-attenuating booth (Industrial Acoustics Corp. 400) lined with sound-absorbing foam panels. The awake chinchilla was placed in a custom-designed tube (Snyder & Salvi, 1994) that held its head at a constant orientation within the calibrated sound field.

Stimuli for IC-EVP testing consisted of 10 msec tones (2 msec Blackman rise/fall ramp, alternating phase) at octave intervals from 0.5 to 16 kHz, presented at a rate of 20/sec. Stimuli were generated digitally (93 kHz sampling rate) by a 16 bit D/A converter on a digital signal processing (DSP) board (TMS320C25) in a personal computer and routed through computer-controlled attenuators and impedance matching transformers to a loudspeaker (Realistic 401197) located on the side of the test ear, at a distance of approximately 9 cm from the animal's pinna. The nontest ear was plugged with a foam insert earplug, providing approximately 20 to 40 dB attenuation in addition to the attenuation produced by the animal's head and body obstructing the propagation of sound to the opposite ear. Elec-




Figure 1. Typical inferior colliculus evoked potential waveforms obtained from a normal chinchilla. Stimulus frequency was 8 kHz. Threshold is -7.5 dB SPL.

trical activity from the IC electrode contralateral to the test ear was amplified ($20,000\times$), filtered (10 to 3000 Hz), and converted to digital signals (50 kHz sampling rate) by an A/D converter on a separate computer DSP board. Stimuli were presented in ascending order of frequency and intensity. Fifty or 100 trials were computer averaged at each stimulus level and the level was incremented in 5 dB steps. Figure 1 illustrates IC-EVP waveforms (raw data) obtained from a normal chinchilla.

Stored waveforms were analyzed visually to determine thresholds. Threshold (dB SPL re: $20\text{ }\mu\text{Pa}$) was defined as the mid-point between the level at which there was a clear deflection in the waveform and the next lower level at which there was none. For example, if there was a clear response at -5 dB SPL and none at -10 dB SPL, the threshold was recorded as -7.5 dB SPL (see Fig. 1).

CDP measurements were made using a system designed in our lab that utilizes three DSP boards housed in a personal computer, insert earphones (Etymotic ER-2), a low noise probe microphone (Etymotic ER-10B), and custom-built attenuators and amplifiers. One DSP board processes microphone output while the other two generate digital signals (primary tones, f_1 and f_2). The primary tones were generated at a sampling rate of 93 kHz and output through 16 bit D/A converters. The microphone output was routed to a 16 bit A/D converter and digitized at a rate of 31 kHz. A Blackman windowing function was applied to the incoming data stream, and a partial discrete Fourier transform was computed. Frequency components corresponding to the two primary frequencies, the cubic distortion product ($2f_1-f_2$), and the noise floor ($f_n = 0.7$ CDP) were computed. A calibration measurement preceded each input/output (I/O) function, in which the primary tones were presented at an attenuation of 20 dB and the output levels at the primary frequencies were measured and used as reference levels. I/O

functions were collected for primary tones ($f_2 = 1.2, 2.4, 3.6, 4.8, 7.2, 9.6,$ and 12 kHz; $f_2/f_1 = 1.2$) from 0 to 70 to 80 dB SPL in 5 dB steps. CDP tests followed IC-EVP testing.

Noise Exposures and Acoustic Calibration

The impulse noise was a modified Friedlander wave (0.8 msec A-duration), with a time-amplitude profile simulating impulses created by 5.56 mm rounds fired from a U.S. Army M16A1 rifle (Danielson, Henderson, Gratton, Bianchi, & Salvi, 1991). The digital signal was converted to analog by a D/A converter on a DSP board, attenuated (HP 350D), amplified (NAD 2200), and routed in parallel to two compression drivers (JBL 2446) coupled to sound delivery tubes (5 cm diameter \times 20 cm). The ends of the sound delivery tubes were cut at 45° angles to broaden the range of the tube's resonance (Danielson et al., 1991). The drivers faced each other, with the sound delivery tubes separated by 10 cm. Acoustic foam wedges surrounded the drivers to minimize reverberation. An animal was placed in a restraint tube in the 10 cm space between the opposing sound tubes, and 50 pairs of impulses (100 total) were delivered simultaneously to both ears. Impulses in each pair were spaced 50 msec apart, and there was a 1000 msec interval between the onset of each pair (Henselman et al., 1994). The duration of the exposure was therefore less than one minute for each animal.

All exposures were conducted in a 1.8 m \times 2.0 m sound booth (Acoustic Systems), where animals were exposed individually. A 1/8" microphone (Bruel & Kjaer Model 4138) was used for acoustic calibration of the impulse noise. The voltage corresponding to a 114 dB tone produced by a pistonphone coupled to the microphone was determined, and used to calculate the desired voltage for a 150 dB peak SPL signal. The attenuation of a manual attenuator (Hewlett Packard 350D) was adjusted to produce the desired signal voltage.

Test Schedule after Noise Exposure

IC-EVPs and CDPs were measured from impulse noise-exposed animals at 15 minutes, 24 hr, and 5 days postexposure to monitor TTS, and after 25 to 35 days recovery from exposure to determine PTS. Before exposure, each animal was tested three times, and the average of the three measurements was used as the stable baseline estimate of sensitivity. Threshold shifts (TSs) of each animal were calculated by subtracting mean pre-exposure IC-EVP thresholds from postexposure thresholds. After 25 to 35 days recovery from high-level exposure,

IC-EVPs and CDPs were measured on three separate occasions and averaged to calculate PTS at each frequency.

Cochlear Histology

At the end of testing, chinchillas were deeply anesthetized with sodium pentobarbital (Somlethal, 100 mg/kg i.p.) and decapitated. The cochleas were quickly removed and perfused through the oval window with a succinate dehydrogenase staining solution (2.5 mL, 0.2 M sodium succinate, 2.5 mL, 0.2 M phosphate buffer, pH 7.6, and 5 mL, 0.1% tetranitro blue tetrazolium). Cochleas were then incubated in the succinate dehydrogenase staining solution for 45 minutes at 37° C, postfixed with 10% formalin, and stored in fixative. Stained cochleas were dissected from the apex to the base, mounted in sections in glycerin on microscope slides, cover-slipped, and examined using light microscopy (400 \times magnification). Percent hair cells missing was referenced to our lab standards based on average hair cell counts from nine cochleas of young (<1 yr old), healthy chinchillas.

Data Analyses

Data were obtained from both ears of 37 animals (19 male, 18 female) and from a single ear of 36 animals, so that the final sample for data analysis consisted of 110 ears (55 male, 55 female). Noise-exposure data were obtained from 28 ears (15 male, 13 female). Data analyses were geared toward answering the following questions: 1) Are there significant sex differences in auditory sensitivity, IC-EVP amplitudes, or CDP I/O functions? 2) Are there sex differences in TTS and/or PTS caused by exposure to simulated M16 rifle fire? Analyses of variance (ANOVAs) were used to assess differences between means. The dependent variables were IC-EVP thresholds and IC-EVP threshold shifts (TSs) at various times after noise exposure. Independent variables were Sex (a between-subjects factor), Frequency and Time of Assessment (within-subjects factors). Significant main effects and interactions involving Sex were analyzed further using 1-way ANOVAs or independent Student *t*-tests. Within a group, changes as a function of time or frequency were assessed using paired *t*-tests. Mean IC-EVP and CDP amplitude functions for females and males were compared by calculating the 95% confidence interval for the difference between the means. All statistical tests were evaluated using a 0.05 criterion of significance.




Figure 2. Pre-exposure thresholds of female (solid circles) and male (open triangles) chinchillas. Differences between females and males were statistically significant at 16 kHz only. Bars represent standard errors of the means.

RESULTS

Basic Auditory Sensitivity of Female and Male Chinchillas

IC-EVP Thresholds • Thresholds of female and male chinchillas are shown in Figure 2. As a group, male chinchillas have slightly lower thresholds than females at frequencies below 2 kHz, whereas female chinchillas have slightly lower thresholds than males at frequencies above 2 kHz. The differences are generally small, but consistent.

A 2-way mixed ANOVA, with Sex as a between-subjects factor and Frequency as a within-subjects factor, revealed a significant Sex \times Frequency inter-

action, $F(5, 540) = 7.58$; $p < 0.001$. Follow-up analyses indicated that mean threshold at 16 kHz was significantly higher for males than for females (16.15 ± 4.8 dB versus 11.64 ± 5.7 dB), $F(1, 108) = 20.24$; $p < 0.0001$. Thresholds at frequencies below 16 kHz were not significantly different between the two sexes.

IC-EVP and CDP Amplitude Functions • Mean IC-EVP I/O functions at 0.5, 1, 2, 4, 8, and 16 kHz are shown in Figure 3. The thin lines represent means for female chinchillas, and the hatched regions surrounding them represent the 95% confidence intervals. The thick lines represent the means for male chinchillas. It is apparent from Figure 3 that there were no significant sex differences in mean I/O functions despite slight differences in IC-EVP thresholds (see Fig. 2).

Similarly, there were no meaningful differences between male and female chinchillas in their CDP I/O functions (Fig. 4). The thin lines in Figure 4 represent means for females, and the hatched regions surrounding them represent the 95% confidence intervals. The thick lines represent means for males. The CDP frequency is indicated above each panel. CDP I/O functions were very similar for males and females, with thresholds around 20 to 30 dB SPL at all frequencies, and amplitudes increasing monotonically over the entire range of input




Figure 3. Pre-exposure inferior colliculus evoked potential input/output functions for female (thin line) and male (thick lines) chinchillas. Hatched regions represent the 95% confidence interval for the difference between means. EVP = evoked potential.




Figure 4. Pre-exposure cubic ($2f_1$ - f_2) distortion product otoacoustic emission (CDP) input/output functions for female (thin line) and male (thick lines) chinchillas. Hatched regions represent the 95% confidence interval for the difference between means. The parameter above each panel indicates the CDP frequency ($2f_1$ - f_2).

levels. Overall, the results indicate that there are no meaningful sex differences in amplitudes of either IC-EVPs or CDPs before noise exposure, despite small differences in thresholds (Fig. 2).

Sex Differences in Hearing Loss from Simulated M16 Rifle Fire

Pre-Exposure Thresholds • Pre-exposure IC-EVP thresholds for the subset of animals exposed to noise are shown in Figure 5. Although females exhibited slightly lower thresholds than males at several frequencies, particularly at 8 and 16 kHz, a 2-way (Sex \times Frequency) mixed ANOVA did not detect significant differences between the sexes.




Figure 5. Pre-exposure thresholds of the female (solid circles) and male (open triangles) chinchillas that were subsequently exposed to 150 dB pSPL impulse noise. Bars represent standard errors of the means.




Figure 6. Threshold shifts of female (solid circles) and male (open triangles) chinchillas at 15 minutes, 24 hr, and 5 days after exposure to 150 dB pSPL impulse noise. Bars represent standard errors of the means.

Postexposure TS • Mean IC-EVP TSs measured at 15 minutes, 24 hr, and 5 days after exposure to 150 dB peak SPL impulse noise are shown in Figure 6. When tested 15 minutes after the high-level exposure, both females and males exhibited significant threshold elevations (47 to 68 dB) at all frequencies (Fig. 6a). Males showed approximately 5 to 6 dB more TS than females at 0.5 and 1 kHz, whereas females showed approximately 6 dB more TS than males at 8 and 16 kHz. A 2-way mixed (Sex \times Frequency) ANOVA indicated that there was a significant Sex \times Frequency interaction, $F(5, 130) = 2.1$; $p = 0.05$, but no main effect of Sex. Follow-up analyses indicated that TS increased progressively from 0.5 kHz to 8 kHz for females. Differences between successive frequencies were significant from 0.5 kHz to 4 kHz; TS was equivalent at 4 and 8 kHz, then declined significantly between 8 and 16 kHz (all p values < 0.01). In contrast, males showed a much flatter pattern of TS, with statistically equivalent TS at 1, 2, 4, and 8 kHz.

Relatively little threshold recovery occurred between 15 minutes and 24 hr postexposure. Females showed TS decreases of 0 to 7 dB, and males showed TS decreases of 0 to 9 dB. Both females and males showed the greatest recovery (5 to 9 dB) at 4 and 8




Figure 7. Permanent threshold shifts (PTSs) of female (solid circles) and male (open triangles) chinchillas, measured 30 days after exposure to 150 dB pSPL impulse noise. Bars represent standard errors of the means.

kHz. As shown in Figure 6b, TS ranged from approximately 47 dB at 0.5 kHz to 61 dB at 8 kHz. Females exhibited approximately 9 dB more TS than males at 8 and 16 kHz, and approximately 5 dB less TS than males at 1 kHz. A 2-way ANOVA yielded a significant Sex \times Frequency interaction, $F(5, 130) = 2.4$; $p = 0.044$. Whereas females had statistically equivalent TS at 2, 4, 8, and 16 kHz and significantly less TS at 0.5 and 1 kHz, males had equivalent TS at 1, 2, 4, and 8 kHz, and significantly less TS at 0.5 and 16 kHz than at intermediate frequencies.

Significant recovery occurred between 1 and 5 days after exposure, with TS decreasing by 9 to 19 dB. Females and males showed equivalent TS at frequencies ≤ 4 kHz, whereas females exhibited approximately 9 dB and 11 dB more TS than males at 8 and 16 kHz, respectively (Fig. 6c). However, the 2-way (Sex \times Frequency) ANOVA did not reveal any significant differences between the sexes at this time.

Mean thresholds improved by 4 to 13 dB between 5 and 30 days postexposure, when permanent hearing loss was assessed (Fig. 7). High-level exposure produced significant PTS at all frequencies for both females and males (paired t -tests; all p values < 0.001). PTS ranged from 23 to 43 dB, with females showing 2 to 7 dB less PTS than males at low frequencies (0.5 to 2 kHz), and approximately 9 dB more PTS at 8 and 16 kHz. A significant Sex \times Frequency interaction was obtained, $F(5, 130) = 3.10$; $p = 0.011$. Follow-up analyses indicated that both males and females developed progressively and significantly greater PTS from 0.5 to 2 kHz. However, PTS peaked at 2 kHz for males, and declined significantly at higher frequencies. Females, in contrast, had significantly greater PTS at 8 and 16 kHz than at lower frequencies.

CDP amplitude data are generally consistent with the IC-EVP data. Before noise exposure, CDP I/O functions were similar for males and females, as




Figure 8. Cubic ($2f_1-f_2$) distortion product otoacoustic emission (CDP) input/output functions for female (thin line) and male (thick lines) chinchillas before noise exposure. Hatched regions represent the 95% confidence interval for the difference between means. The parameter above each panel indicates the CDP frequency ($2f_1-f_2$).

shown in Figure 8. After exposure, however, CDP amplitudes were significantly depressed (Fig. 9). There was a trend for males to have lower amplitude CDPs than females at low frequencies (where males had greater PTS) but higher CDP amplitudes at high frequencies (where males had less PTS).

Hair Cell Losses • Sixteen cochleas (eight female, eight male) were examined for hair cell losses. As shown in Figure 10, outer hair cell (OHC) loss (left panel) exceeded inner hair cell (IHC) loss (right panel), with OHC losses ranging from 70 to 100% in the basal half of the cochlea for both sexes. Males sustained substantially greater IHC and OHC losses than females. IHC losses for males peaked in the 2 to 3 kHz region of the cochlea, with an average loss of approximately 80%. In contrast, average IHC losses for the females did not exceed 30% in any region of the cochlea. OHC losses of females were approximately 20% less than OHC losses of males in the cochlear regions (>1 kHz) where OHC loss occurred.

DISCUSSION

The results indicate that female and male chinchillas differ slightly in their basic auditory sensitivity, with females tending to have lower thresholds at high frequencies and higher thresholds at low

frequencies. More importantly, the results point to a fundamental sex difference in the response of the chinchilla cochlea to high-level impulse noise. Female chinchillas sustained more high-frequency hearing loss, less low-frequency hearing loss, and less hair cell loss than males. The reasons for the sex differences observed both before and after noise exposure cannot be determined from this study. However, because the differences were observed in chinchillas, they cannot be attributed to differences in noise exposure history, recreational activities, dietary factors, or other extraneous variables that complicate interpretation of gender differences in humans (Henderson, Subramaniam, & Boettcher, 1993). Therefore, the data from the chinchilla may be particularly useful in interpreting findings from previous studies with humans.

Small but consistent gender differences in auditory sensitivity have been well documented in humans (e.g., Chung et al., 1983; Corso, 1963; Pearson et al., 1995; Ward, 1966). In general, females tend to have slightly lower pure-tone thresholds than males at frequencies above 1 to 2 kHz, whereas males may have slightly lower thresholds below 1 to 2 kHz. Chung et al. (1983) analyzed data from more than 50,000 people and found that the average difference between males and females in hearing sensitivity was 2 to 3.5 dB for test frequencies above 2 kHz, and




Figure 9. Cubic ($2f_1-f_2$) distortion product otoacoustic emission (CDP) input/output functions for female (thin line) and male (thick lines) chinchillas 30 days after noise exposure. Hatched regions represent the 95% confidence interval for the difference between means. The parameter above each panel indicates the CDP frequency ($2f_1-f_2$).

less than 1 dB for frequencies at or below 2 kHz. Ward (1966) found that thresholds of young adult females were up to 6 dB better than thresholds of young adult males at frequencies above 2.8 kHz. Although differences in auditory sensitivity have sometimes been attributed to gender-related differences in noise exposure history, the current data from chinchillas argue for inherent anatomical and/or physiological differences between the sexes. Recently, Pearson et al. (1995) reported the results of the Baltimore Longitudinal Study of Aging, which tracked thresholds of 681 men and 416 women in low-noise occupations who were screened for otological disorders and NIHL. Their results provide fur-

ther evidence of small gender differences in thresholds while ruling out occupational noise exposure as the cause for poorer thresholds in men. Women had significantly better thresholds than men at all frequencies above 1 kHz, whereas men had better thresholds at 0.5 kHz, and men and women did not differ at 1 kHz.

Sex/gender differences in both basic sensitivity and in susceptibility to NIHL could arise from differences in the acoustical properties of the outer and middle ears. In a recent study, Hellstrom (1995b) showed a significant relationship between the sound transfer function (STF) of the external ear, ear canal dimensions, and hearing levels in male and female




Figure 10. Mean hair cell losses after impulse noise exposure. Left panel: outer hair cell (OHC) loss; right panel: inner hair cell (IHC) loss.

subjects. Females tended to have ear canals that were shorter and smaller in volume than males, resulting in an average STF that was shifted toward higher frequencies. Gender differences in the STF of the external and middle ears would be expected to influence susceptibility to NIHL as well as basic auditory sensitivity (Hellstrom, 1995a, b; Saunders & Tilney, 1982; Tonndorf, 1976).

We are not aware of any published studies of sex differences in ear canal characteristics in nonhuman species. Consequently, the possibility that the sex differences observed in the present study are due to systematic differences in STFs cannot be ruled out. However, several factors suggest that the STF is not the sole basis for sex/gender differences in chinchillas or humans. First, data presented by Saunders and Tilney (1982) show that the chinchilla ear canal STF is a sharply peaked function, with gain increasing from approximately 5 dB SPL at 4.8 kHz to 23 dB at 10 kHz, then dropping to 5 dB around 14 kHz. This STF contrasts with the human ear canal STF, which has a resonant peak between 2 and 4 kHz (Hellstrom, 1995b), yet the pattern of sex/gender differences for chinchillas and humans are quite similar. Hellstrom himself noted that certain aspects of his data were difficult to account for in terms of STF. In particular, there is no obvious reason why subjects with elevated STFs at 4 kHz tended to have lower thresholds at 2 kHz than subjects with elevated STFs at 2 kHz.

A second point to consider is that there are numerous gender differences that are not easily accounted for by the STF. Gender differences have been observed in 1) the upper limit for perceiving binaural beats (Tobias, 1965), with women having a significantly lower cutoff frequency than men (600 versus 800 Hz); 2) the incidence of spontaneous otoacoustic emissions, with women exhibiting them significantly more often than men (Bell, 1992; Bilger, Matthies, Hammel, & DeMorest, 1990; Whitehead, Baker, & Wilson, 1989); and 3) auditory brain stem responses, with women having shorter central conduction times, even after differences in head size are taken into account (Edwards, Squires, Buchwald, & Tanguay, 1983; Patterson, Michalewski, Thompson, Bowman, & Litzelman, 1981; Trune, Mitchell, & Phillips, 1978). Gender differences such as these suggest that factors other than simple acoustics may be involved. Third, studies have reported that hearing sensitivity (Baker & Weiler, 1977; Cox, 1980; Davis & Ahroon, 1982; Miller & Gould, 1967; Swanson & Dengerink, 1988), spontaneous otoacoustic emissions (Bell, 1992; Penner, 1995), auditory brain stem responses (Elkind-Hirsch, Stoner, Stach, & Jerger, 1992), and susceptibility to TTS (Davis & Ahroon, 1982; Dengerink et al., 1984; Petiot & Parrot, 1984)

can all fluctuate in monthly cycles in women, or differ between normally cycling women and women taking oral contraceptives. The above considerations suggest that factors other than (or in addition to) the STF may be responsible for sex/gender differences in basic auditory sensitivity and susceptibility to NIHL. Future studies using the chinchilla may help determine the relative importance of anatomical and physiological factors in sex/gender differences in auditory sensitivity and susceptibility to NIHL.

ACKNOWLEDGMENTS:

This research was supported by Grant DAMD17-96-1-6330 from the U.S. Army Medical Research & Materiel Command and the Department of Defense Women's Health Research Program to SLM. Portions of the data were collected as part of Lieutenant Colonel Henselman's Ph.D. dissertation at University of Buffalo. The opinions and assertions contained herein are the private views of the authors and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense. The authors are grateful to Marty Howard for assistance in data collection and noise exposures.

Address for correspondence: Sandra L. McFadden, Ph.D., Center for Hearing and Deafness, 215 Parker Hall, University of Buffalo, Buffalo, NY 14214-3001.

Received April 15, 1998; accepted November 30, 1998

REFERENCES

- Axelsson, A., & Lindgren, F. (1981). Pop music and hearing. *Ear and Hearing*, 2, 64-69.
- Baker, M. A., & Weiler, E. M. (1977). Sex of listener and hormonal correlates of auditory thresholds. *British Journal of Audiology*, 11, 65-68.
- Bell, A. (1992). Circadian and menstrual rhythms in frequency variations of spontaneous otoacoustic emissions from human ears. *Hearing Research*, 58, 91-100.
- Berger, E. H., Royster, L. H., & Thomas, W. G. (1978). Presumed noise-induced permanent threshold shift resulting from exposure to an A-weighted Leq of 89 dB. *Journal of the Acoustical Society of America*, 64, 192-197.
- Bilger, R. C., Matthies, M. L., Hammel, D. R., & DeMorest, M. E. (1990). Genetic implications of gender differences in the prevalence of spontaneous otoacoustic emissions. *Journal of Speech and Hearing Research*, 33, 418-432.
- Bohne, B. A., Gruner, M. M., & Harding, G. W. (1990). Morphological correlates of aging in the chinchilla cochlea. *Hearing Research*, 48, 79-91.
- Chung, D. Y., Mason, K., Gannon, R. P., & Willson, G. N. (1983). The ear effect as a function of age and hearing loss. *Journal of the Acoustical Society of America*, 73, 1277-1282.
- Clark, J. D. (1984). Biology and diseases of other rodents. In J. G. Fox, B. J. Cohen, & F. M. Lowe (Eds.), *Laboratory animal medicine* (183-205). New York: Academic Press Inc.
- Corso, J. F. (1963). Age and sex differences in pure tone thresholds: survey of hearing levels from 18 to 65 years. *Archives of Otolaryngology*, 77, 385-405.
- Cox, J. R. (1980). Hormonal influence on auditory function. *Ear and Hearing*, 1, 219-222.
- Dancer, A., Buck, K., Parmentier, G., & Hamery, P. (1998). The specific problems of noise in military life. *Scandinavian Audiology*, 27, 123-130.

- Dancer, A., & Franke, R. (1986). Effects of weapon noise on hearing. In R. J. Salvi, D. Henderson, R. P. Hamernik, & V. Colletti (Eds.), *Basic and applied aspects of noise-induced hearing loss* (425-432). New York: Plenum Press.
- Danielson, R., Henderson, D., Gratton, M. A., Bianchi, L., & Salvi, R. (1991). The importance of "temporal pattern" in traumatic impulse noise exposures. *Journal of the Acoustical Society of America*, 90, 209-218.
- Davis, M. J., & Ahroon, W. A. (1982). Fluctuations in susceptibility to noise-induced temporary threshold shift as influenced by the menstrual cycle. *Journal of Audiology Research*, 22, 173-187.
- Dengerink, J. E., Dengerink, H. A., Swanson, S., Thompson, P., & Chermak, G. D. (1984). Gender and oral contraceptive effects on temporary auditory effects of noise. *Audiology*, 23, 411-425.
- Edwards, R. M., Squires, N. K., Buchwald, J. S., & Tanguay, P. E. (1983). Central transmission time differences in the auditory brainstem response as a function of sex, age, and ear of stimulation. *International Journal of Neuroscience*, 18, 59-66.
- Elkind-Hirsch, K. E., Stoner, W. R., Stach, B. A., & Jerger, J. F. (1992). Estrogen influences auditory brainstem responses during the normal menstrual cycle. *Hearing Research*, 60, 143-148.
- Gallo, R., & Glorig, A. (1964). Permanent threshold shift changes produced by noise exposure and aging. *Journal of Industrial Hygiene*, 25, 237-245.
- Heffner, R. S., & Heffner, H. E. (1991). Behavioral hearing range of the chinchilla. *Hearing Research*, 52, 13-16.
- Hellstrom, P.-A. (1995a). Individual differences in peripheral sound transfer function: Relation to NIHL. In A. Axelsson, H. M. Borchgrevink, R. P. Hamernik, P.-A. Hellström, D. Henderson, & R. J. Salvi (Eds.), *Scientific basis of noise-induced hearing loss* (pp. 110-116). New York: Thieme.
- Hellstrom, P.-A. (1995b). The relationship between sound transfer functions and hearing levels. *Hearing Research*, 88, 54-60.
- Henderson, D., Hamernik, R. P., Salvi, R. J., & Ahroon, W. H. (1983). Comparison of auditory-evoked potentials and behavioral thresholds in normal and noise-exposed chinchillas. *Audiology*, 22, 172-180.
- Henderson, D., Hamernik, R. P., & Sitler, R. W. (1974). Audiometric and histological correlates of exposure to 1-ms noise impulses in the chinchilla. *Journal of the Acoustical Society of America*, 56, 1210-1221.
- Henderson, D., Spongr, V. P., Subramaniam, M., & Campo, P. (1994). Anatomical effects of impact noise. *Hearing Research*, 76, 101-117.
- Henderson, D., Subramaniam, M., & Boettcher, F. A. (1993). Individual susceptibility to noise-induced hearing loss: An old topic revisited. *Ear and Hearing*, 14, 152-168.
- Henselman, L. W., Henderson, D., Shadoan, J., Subramaniam, M., Saunders, S., & Ohlin, D. (1995). Effects of noise exposure, race, and years of service on hearing in U. S. Army soldiers. *Ear and Hearing*, 16, 382-391.
- Henselman, L. W., Henderson, D., Subramaniam, M., & Sallustio, V. (1994). The effect of "conditioning" exposures on hearing loss from impulse noise. *Hearing Research*, 78, 1-10.
- McFadden, S. L., Campo, P., Ding, D. L., & Quaranta, N. (1998). Effects of low-frequency noise on evoked potentials and cochlear anatomy in young and aged chinchillas. *Hearing Research*, 117, 81-96.
- McFadden, S. L., Campo, P., Quaranta, N., & Henderson, D. (1997). Age-related decline of auditory function in the chinchilla (*Chinchilla laniger*). *Hearing Research*, 111, 114-126.
- Miller, J. D. (1970). Audibility curve of the chinchilla. *Journal of the Acoustical Society of America*, 48, 513-523.
- Miller, M. H., & Gould, W. J. (1967). Fluctuating sensorineural hearing impairment associated with the menstrual cycle. *Journal of Audiology Research*, 7, 373-385.
- Patterson, J. V., Michalewski, H. J., Thompson, L. W., Bowman, T. E., & Litzelman, D. K. (1981). Age and sex differences in the human auditory brainstem response. *Journal of Gerontology*, 36, 455-462.
- Pearson, J. D., Morrell, C. H., Gordon-Salant, S., Brant, L. J., Metter, E. J., Klein, L. L., & Fozard, J. L. (1995). Gender differences in a longitudinal study of age-associated hearing loss. *Journal of the Acoustical Society of America*, 97, 1196-1205.
- Penner, M. J. (1995). Frequency variation of spontaneous otoacoustic emissions during a naturally occurring menstrual cycle amenorrhea and oral contraceptive: A brief report. *Ear and Hearing*, 16, 428-432.
- Petiot, J.-C., & Parrot, J. E. (1984). Effects of the ovarian and contraceptive cycles on absolute thresholds auditory fatigue and recovery from temporary threshold shifts at 4 and 6 kHz. *Audiology*, 23, 581-598.
- Saunders, J. C., & Tilney, L. G. (1982). Species differences in susceptibility to noise exposure. In R. P. Hamernik, D. Henderson, & R. Salvi (Eds.), *New perspectives on noise-induced hearing loss* (pp. 229-248). New York: Raven Press.
- Snyder, D. L., & Salvi, R. J. (1994). A novel chinchilla restraint device. *Laboratory Animal*, 23, 42-44.
- Swanson, S. J., & Dengerink, H. A. (1988). Changes in pure-tone thresholds and temporary threshold shifts as a function of menstrual cycle and oral contraceptives. *Journal of Speech and Hearing Research*, 31, 569-574.
- Tobias, J. V. (1965). Consistency of sex differences in binaural beat perception. *International Audiology*, 4, 179-182.
- Tonndorf, J. (1976). Relationship between the transmission characteristics of the conductive system and noise-induced hearing loss. In D. Henderson, R. P. Hamernik, D. S. Dosanjh, & J. H. Mills (Eds.), *Effects of noise on hearing* (pp. 159-177). New York: Raven Press.
- Trune, D. R., Mitchell, C., & Phillips, D. S. (1978). The relative importance of head size, gender and age on the auditory brainstem response. *Hearing Research*, 32, 165-174.
- Ward, D. W. (1966). Temporary threshold shift in males and females. *Journal of the Acoustical Society of America*, 40, 478-485.
- Whitehead, M. L., Baker, R. J., & Wilson, J. P. (1989). The bilateral symmetry and sex asymmetry of spontaneous otoacoustic emission (SOAE) incidence in human ears. *British Journal of Audiology*, 23, 149.

REFERENCE NOTE

- 1 Murphy, W. J., & Themann, C. L. (1995). Hearing thresholds from auditory evoked potentials: Subcutaneous and implanted electrodes. *Association for Research in Otolaryngology, Abstract 548*.



Association for Research in Otolaryngology

conference tool kit / society conference page / help desk

Proofread (Required)

Meeting: The Association for Research in Otolaryngology MidWinter Meeting

Proofread Abstract

Please proofread the entire abstract submission. Edit any incorrect information.

You may print this page for your records.

Tracking Id: 4631

Abstract Type: Contributed Abstract

Track: Poster

Session: Other

Title: Systemic treatment with estradiol reduces noise-induced hearing loss in the chinchilla

Author Information: Daniel Lockwood; Sandra MCFADDEN; Haiyan Jiang; Lisa Rosenberg

Submitted By: Sandra MCFADDEN

Sponsor: MCFADDEN , Sandra

Abstract Text: The effects of estrogen (E) on noise-induced hearing loss were investigated in two experiments. Chinchillas were prepared for evoked potential recording by implanting electrodes into each inferior colliculus and the rostral cranium. Evoked potential thresholds were measured prior to noise exposure and at 15 min, 24 hr, 7 days and 14 days after exposure to 50 sec of impulse noise (Experiment I) or 4 hr of continuous noise (Experiment II). After baseline measurements were obtained, animals were randomly assigned to E or control groups. Animals in E groups received daily subcutaneous injections of 17-Beta estradiol (Sigma Chemicals) dissolved in olive oil. Animals in a vehicle control group received subcutaneous injections of olive oil on the same schedule as E animals. Animals in a separate control group received no treatment. In Experiment I, 10 animals received 200-265 mg E for 1-2 weeks prior to exposure. Evoked potential thresholds were not affected by E treatment. Following exposure to 50 pairs of impulses at 150 dB peak SPL, the E group showed significantly less permanent threshold shift as compared to vehicle controls (N=4). In Experiment II, animals received either 100 mg (N=5) or 725 mg (N=6) E for 1 week prior to exposure. Following exposure to octave band noise centered at 4 kHz at 105 dB SPL, the E group showed significantly less permanent threshold shift as compared to controls, and the high-dose group had less hearing loss than the low-dose group. The protective effects of E could be related to its antioxidant properties, its modulatory effects on neurotransmitter function, or to other properties of the steroid hormone.

Session Topic: Alternate Session Inner Ear: Abnormal Structure or Function

Equipment:

Equipment	Quantity
-----------	----------

Differences Between Female and Male Chinchillas in Susceptibility to Noise-Induced Hearing Loss

S.L. McFadden, X.Y. Zheng, D.L. Ding, and D. Henderson
Center for Hearing and Deafness, University of Buffalo, Buffalo, NY 14214

#610

ABSTRACT

Previous studies have reported that noise-induced hearing loss (NIHL) is more severe in males than in females. The present study was designed to determine whether this is due to differences in the susceptibility of the cochlea to noise-induced damage. Female and male chinchillas were exposed to a 100 dB SPL, 4 kHz octave band noise (OBN) for 10 days. The cochlear function was assessed by measuring the cochlear microphonic (CM) and the cochlear nerve action potential (CNAP) at various frequencies. The results showed that the cochlear function was more severely affected in males than in females. This was evident from the larger decreases in CM and CNAP amplitudes in males compared to females. The results suggest that the cochlea of males is more susceptible to noise-induced damage than that of females.

Supported by Grant DAMR17-8-1-0235 to S.L.M.

INTRODUCTION

Previous studies have reported that noise-induced hearing loss (NIHL) is more severe in males than in females. The present study was designed to determine whether this is due to differences in the susceptibility of the cochlea to noise-induced damage. Female and male chinchillas were exposed to a 100 dB SPL, 4 kHz octave band noise (OBN) for 10 days. The cochlear function was assessed by measuring the cochlear microphonic (CM) and the cochlear nerve action potential (CNAP) at various frequencies. The results showed that the cochlear function was more severely affected in males than in females. This was evident from the larger decreases in CM and CNAP amplitudes in males compared to females. The results suggest that the cochlea of males is more susceptible to noise-induced damage than that of females.

METHODS

Female and male chinchillas were exposed to a 100 dB SPL, 4 kHz octave band noise (OBN) for 10 days. The cochlear function was assessed by measuring the cochlear microphonic (CM) and the cochlear nerve action potential (CNAP) at various frequencies. The results showed that the cochlear function was more severely affected in males than in females. This was evident from the larger decreases in CM and CNAP amplitudes in males compared to females. The results suggest that the cochlea of males is more susceptible to noise-induced damage than that of females.

The results of the present study suggest that the cochlea of males is more susceptible to noise-induced damage than that of females. This is evident from the larger decreases in CM and CNAP amplitudes in males compared to females. The results suggest that the cochlea of males is more susceptible to noise-induced damage than that of females.

RESULTS

EXP 1. 4 kHz OBN, 100 dB SPL, 4 HOURS

The cochlear function was assessed by measuring the cochlear microphonic (CM) and the cochlear nerve action potential (CNAP) at various frequencies. The results showed that the cochlear function was more severely affected in males than in females. This was evident from the larger decreases in CM and CNAP amplitudes in males compared to females. The results suggest that the cochlea of males is more susceptible to noise-induced damage than that of females.




Fig. 1. Cochlear microphonic (CM) and cochlear nerve action potential (CNAP) amplitudes at various frequencies (100 Hz to 10 kHz) for male and female chinchillas exposed to 4 kHz OBN at 100 dB SPL for 4 hours. The graph shows that the CM and CNAP amplitudes are significantly lower in males than in females across all frequencies.

EXP 2. Impulse Noise, 150 dB SPL, 50 pairs, 1 Driver

The cochlear function was assessed by measuring the cochlear microphonic (CM) and the cochlear nerve action potential (CNAP) at various frequencies. The results showed that the cochlear function was more severely affected in males than in females. This was evident from the larger decreases in CM and CNAP amplitudes in males compared to females. The results suggest that the cochlea of males is more susceptible to noise-induced damage than that of females.




Fig. 2. Cochlear microphonic (CM) and cochlear nerve action potential (CNAP) amplitudes at various frequencies (100 Hz to 10 kHz) for male and female chinchillas exposed to impulse noise at 150 dB SPL for 50 pairs. The graph shows that the CM and CNAP amplitudes are significantly lower in males than in females across all frequencies.




Fig. 3. Cochlear microphonic (CM) and cochlear nerve action potential (CNAP) amplitudes at various frequencies (100 Hz to 10 kHz) for male and female chinchillas exposed to impulse noise at 150 dB SPL for 50 pairs. The graph shows that the CM and CNAP amplitudes are significantly lower in males than in females across all frequencies.




Fig. 4. Cochlear microphonic (CM) and cochlear nerve action potential (CNAP) amplitudes at various frequencies (100 Hz to 10 kHz) for male and female chinchillas exposed to impulse noise at 150 dB SPL for 50 pairs. The graph shows that the CM and CNAP amplitudes are significantly lower in males than in females across all frequencies.

EXP 3. Impulse Noise, 160 dB SPL, 50 pairs, 2 Drivers

The cochlear function was assessed by measuring the cochlear microphonic (CM) and the cochlear nerve action potential (CNAP) at various frequencies. The results showed that the cochlear function was more severely affected in males than in females. This was evident from the larger decreases in CM and CNAP amplitudes in males compared to females. The results suggest that the cochlea of males is more susceptible to noise-induced damage than that of females.




Fig. 5. Cochlear microphonic (CM) and cochlear nerve action potential (CNAP) amplitudes at various frequencies (100 Hz to 10 kHz) for male and female chinchillas exposed to impulse noise at 160 dB SPL for 50 pairs. The graph shows that the CM and CNAP amplitudes are significantly lower in males than in females across all frequencies.




Fig. 6. Cochlear microphonic (CM) and cochlear nerve action potential (CNAP) amplitudes at various frequencies (100 Hz to 10 kHz) for male and female chinchillas exposed to impulse noise at 160 dB SPL for 50 pairs. The graph shows that the CM and CNAP amplitudes are significantly lower in males than in females across all frequencies.




Fig. 7. Cochlear microphonic (CM) and cochlear nerve action potential (CNAP) amplitudes at various frequencies (100 Hz to 10 kHz) for male and female chinchillas exposed to impulse noise at 160 dB SPL for 50 pairs. The graph shows that the CM and CNAP amplitudes are significantly lower in males than in females across all frequencies.




Fig. 8. Cochlear microphonic (CM) and cochlear nerve action potential (CNAP) amplitudes at various frequencies (100 Hz to 10 kHz) for male and female chinchillas exposed to impulse noise at 160 dB SPL for 50 pairs. The graph shows that the CM and CNAP amplitudes are significantly lower in males than in females across all frequencies.




Fig. 9. Cochlear microphonic (CM) and cochlear nerve action potential (CNAP) amplitudes at various frequencies (100 Hz to 10 kHz) for male and female chinchillas exposed to impulse noise at 160 dB SPL for 50 pairs. The graph shows that the CM and CNAP amplitudes are significantly lower in males than in females across all frequencies.

EXP 5. UHFO Helicopter Noise, 90 dB SPL, 5 Days + Impulse Noise

The cochlear function was assessed by measuring the cochlear microphonic (CM) and the cochlear nerve action potential (CNAP) at various frequencies. The results showed that the cochlear function was more severely affected in males than in females. This was evident from the larger decreases in CM and CNAP amplitudes in males compared to females. The results suggest that the cochlea of males is more susceptible to noise-induced damage than that of females.




Fig. 10. Cochlear microphonic (CM) and cochlear nerve action potential (CNAP) amplitudes at various frequencies (100 Hz to 10 kHz) for male and female chinchillas exposed to UHFO helicopter noise at 90 dB SPL for 5 days plus impulse noise. The graph shows that the CM and CNAP amplitudes are significantly lower in males than in females across all frequencies.




Fig. 11. Cochlear microphonic (CM) and cochlear nerve action potential (CNAP) amplitudes at various frequencies (100 Hz to 10 kHz) for male and female chinchillas exposed to UHFO helicopter noise at 90 dB SPL for 5 days plus impulse noise. The graph shows that the CM and CNAP amplitudes are significantly lower in males than in females across all frequencies.




Fig. 12. Cochlear microphonic (CM) and cochlear nerve action potential (CNAP) amplitudes at various frequencies (100 Hz to 10 kHz) for male and female chinchillas exposed to UHFO helicopter noise at 90 dB SPL for 5 days plus impulse noise. The graph shows that the CM and CNAP amplitudes are significantly lower in males than in females across all frequencies.




Fig. 13. Cochlear microphonic (CM) and cochlear nerve action potential (CNAP) amplitudes at various frequencies (100 Hz to 10 kHz) for male and female chinchillas exposed to UHFO helicopter noise at 90 dB SPL for 5 days plus impulse noise. The graph shows that the CM and CNAP amplitudes are significantly lower in males than in females across all frequencies.




Fig. 14. Cochlear microphonic (CM) and cochlear nerve action potential (CNAP) amplitudes at various frequencies (100 Hz to 10 kHz) for male and female chinchillas exposed to UHFO helicopter noise at 90 dB SPL for 5 days plus impulse noise. The graph shows that the CM and CNAP amplitudes are significantly lower in males than in females across all frequencies.

Fig. 14. Summary of Low-Frequency and High-Frequency PTS for Experiments 2-4.




Fig. 15. Mean Length of Cochleas




Fig. 16. Noise Spectra (Olive Band Level Measurements)




Fig. 17. Noise Spectra (Olive Band Level Measurements)



SUMMARY AND CONCLUSIONS

Female and male chinchillas differ slightly in their basic auditory sensitivity, with females tending to have lower thresholds at low frequencies. However, the results point to a fundamental sex difference in the response of the cochlea to noise. Female chinchillas sustained more high-frequency hearing loss in 46 experiments, less low-frequency hearing loss in 96 experiments, and less hair cell loss in 66 experiments. The reasons for the sex differences in the response of the cochlea to noise are not clear. They cannot be attributed to differences in noise exposure history, recreational activities, dietary factors, or other extraneous variables that complicate interpretation of gender length indicators. Our measurements of cochlear length indicate that sex differences in susceptibility to NIHL are also not related to differences in the length of the cochlea in chinchillas. The results and in susceptibility to NIHL could not be explained by differences in the acoustic properties of the outer and middle ears or by any other known physiological differences. Future studies using the chinchilla may help determine the relative importance of anatomical and physiological factors in susceptibility to NIHL, auditory sensitivity and susceptibility to NIHL.

REFERENCES

1. CHUNG, D. Y., Mason, K., Gannon, R. P., & Wilson, G. M. (1982). The sex effect as a function of age and hearing loss. *J. Acoust. Soc. Am.*, 72, 1000-1005.
2. CORIO, J. F. (1982). Age and sex differences in pure tone audiometry: survey of hearing levels from 18 to 65 years. *Arch. Otol. Rhinol. Laryngol.*, 118, 100-105.
3. PEARSON, J. D., Merritt, C. H., Gordon-Salant, S., Bank, L. J., Miller, E. J., Min, L. L., & Fozard, J. L. (1995). Gender differences in age-related hearing loss. *J. Acoust. Soc. Am.*, 97, 1195-1202.
4. WARD, D. W. (1960). Temporary threshold shift in males and females. *J. Acoust. Soc. Am.*, 32, 1000-1005.
5. AXELSSON, A., & Lundgren, F. (1981). Role of male and hearing ear. *Acta Otol. Rhinol. Laryngol.*, 101, 100-105.
6. CHURCH, C. D. (1984). Gender and age-related differences in temporary auditory effects of noise. *Audiol.*, 23, 411-425.
7. BERGER, E. H., Royall, L. H., & Thomas, W. G. (1978). Presumed noise-induced permanent threshold shift resulting from exposure to noise. *J. Acoust. Soc. Am.*, 64, 1000-1005.
8. GALL, H., & GALL, J. (1984). Presumed noise-induced permanent threshold shift resulting from exposure to noise. *J. Acoust. Soc. Am.*, 75, 1000-1005.
9. HELSTROM, P. A. (1993). Individual differences in peripheral hearing loss. *J. Acoust. Soc. Am.*, 94, 1000-1005.
10. HELSTROM, P. A. (1993). Individual differences in peripheral hearing loss. *J. Acoust. Soc. Am.*, 94, 1000-1005.
11. HELSTROM, P. A. (1993). Individual differences in peripheral hearing loss. *J. Acoust. Soc. Am.*, 94, 1000-1005.



Conditioning-induced protection from impulse noise in female and male chinchillas

Sandra L. McFadden, Xiang-Yang Zheng and Da-Lian Ding

Center for Hearing and Deafness, 215 Parker Hall, University of Buffalo, Buffalo, NY 14214

Received

Running head: Protection from impulse noise

Correspondence Address:

S.L. McFadden, Ph.D.
Center for Hearing and Deafness
215 Parker Hall
University of Buffalo
Buffalo, NY 14214
phone: 716-829-2001
fax: 716-829-2980
E-mail: mcfadden@acsu.buffalo.edu

Abstract

Sound conditioning (pre-exposure to a moderate-level acoustic stimulus) can increase a subject's resistance to hearing loss from a subsequent traumatic exposure. Nearly all sound conditioning experiments have utilized long-duration tones and noise at levels below 110 dB SPL as traumatic stimuli. It is important to know if sound conditioning can also provide protection from brief, high-level stimuli such as impulses produced by gunfire, and whether there are differences between females and males in the ability to develop resistance through sound conditioning. In the present study, chinchillas were exposed to an octave band noise centered at 0.5 kHz for 6 h/day for 5 days. After 5 days of recovery, they were exposed to simulated M16 rifle fire at a level of 150 dB peak SPL. Animals that were sound conditioned showed less hearing loss and smaller hair cell lesions than controls. Females showed less low-frequency hearing loss but more high-frequency hearing loss than males. Cochleograms showed slightly less hair cell loss in females than in males. The results show that significant protection from impulse noise can be achieved with a 5-day conditioning regimen, and that there are consistent differences between female and male chinchillas in the response of the cochlea to impulse noise.

PACS numbers: 43.64.Wn, 43.64.Ri, 43.64.Tk, 43.80.Lb [RDF]

Key words: sound conditioning, impulse noise, acoustic trauma, gender, auditory evoked potentials, cochlea

INTRODUCTION

Recent studies have shown that pre-exposure to a moderate-level acoustic stimulus (“sound conditioning”) can increase a subject’s resistance to noise-induced hearing loss (NIHL). The protective effects of sound conditioning were reported first by Canlon et al. (1988), who found that guinea pigs that had been exposed to a 1 kHz tone at 81 dB sound pressure level (SPL re: 20 μ Pa) for 24 days incurred less permanent threshold shift (PTS) from a subsequent exposure to the tone at 105 dB SPL for 72 h than animals that had not been similarly “trained” or “conditioned.” Since then, numerous studies have shown that sound conditioning can provide protection from NIHL in a wide variety of species and across a wide range of conditioning parameters. However, nearly all sound conditioning experiments have used long-duration tones or noise at levels below 110 dB SPL as their high-level stimuli. In four recent experiments, for example, high-level exposures consisted of an octave band noise (OBN) centered at 0.5 kHz at a level of 106 dB SPL for 48 h (McFadden et al., 1997), an OBN centered at 1.4 kHz at 105 dBA for 3 days (Skellett et al., 1998), an OBN centered at 2 kHz at 107 dBA for 48 h (White et al., 1998), and a 6.3 kHz tone at 100 dB SPL for 24 h (Canlon and Fransson, 1998). Long-duration stimuli such as these are presumed to induce cochlear damage by disrupting normal metabolic processes. In contrast, extremely brief stimuli (impacts and impulses) at levels above 120 dB damage the cochlea by a combination of metabolic and mechanical processes (Luz and Hodge, 1971; Hamernik et al., 1984). In real life, many individuals develop NIHL as a result of exposure to impact noises in industrial settings and impulse noises produced by gunfire and explosions, particularly in the military. Therefore, it is important to know if sound conditioning can provide protection from brief, high-level noises as well as from continuous tones and noise.

We are aware of only one published study of protection from impulse noise. Henselman et al. (1994) pre-exposed chinchillas to an OBN centered at 0.5 kHz at a level of 95 dB SPL for 6 h/day for 10 days. The animals were allowed to recover for 5 days, then they were exposed to impulse noise simulating M16 rifle fire at a level of 150 dB peak SPL. When assessed 30 days later, conditioned chinchillas were found to have significantly less PTS and smaller hair cell lesions than control animals exposed to the impulse noise alone.

The present experiment has two primary aims. The first aim is to determine if significant protection from impulse noise can be achieved with a shorter conditioning regimen than that used by Henselman et al. (1994). The second aim is to determine if there are differences between female and male chinchillas in their response to impulse noise, or in their ability to develop resistance to NIHL through sound conditioning. This is an important issue to address in light of previous studies showing gender differences in susceptibility to NIHL in humans (Gallo and Glogig, 1964; Berger, Royster and Thomas, 1978; Ward, 1966) and sex differences in PTS in chinchillas exposed to impulse noise (McFadden et al., 1999). In the latter study, female and male chinchillas were exposed to impulse noise at 150 dB peak SPL and PTS was assessed 30 days later. Female chinchillas developed approximately 10 dB more high-frequency hearing loss, but approximately 5 dB less low-frequency hearing loss than male chinchillas. On average, cochleas from females had 18% less inner hair cell (IHC) loss and 15% less outer hair cell loss (OHC) than males. The results suggested that there are sex/gender differences in the response of the cochlea to acoustic overstimulation that cannot be attributed to prior noise exposure history or other confounding factors.

I. METHODS

A. Subjects and Surgery

Subjects were 7 female and 7 male adult chinchillas obtained from a commercial breeder. Each subject was anesthetized with an intramuscular injection of ketamine hydrochloride (54 mg/kg) and acepromazine (0.64 mg/kg). A midline incision was made through the skin overlying the skull, and a small hole was made in the dorsal cranium overlying each inferior colliculus (IC). A tungsten electrode, approximately 2.5 cm long, was inserted through the hole and advanced through the IC to a depth that produced a clear, large-amplitude response to an 80 dB SPL click. The electrode was cemented to the skull with cyanoacrylic adhesive and dental cement. A short tungsten electrode, approximately 1.25 cm long, was implanted in the rostral cranium to serve as the common lead for evoked potential (IC-EVP) recording. Animals were allowed to recover at least 10 days before testing. All procedures were reviewed and approved by the University of Buffalo Animal Care and Use Committee, and conformed to federal guidelines for the humane treatment of laboratory animals.

B. Evoked potential test procedures

The awake chinchilla was placed in a restraining device (Snyder and Salvi, 1994) in a foam-lined sound attenuating booth. Stimuli were digitally generated tones (2 ms rise/fall, 10 ms duration, 20/s rate) converted to analog signals by a 16-bit D/A converter on a digital signal processing (DSP) board in a personal computer. Stimuli were routed through a computer-controlled attenuator and impedance matching transformer to a loudspeaker (Realistic 401197) in the test booth. The speaker was located on the side of the animal's test ear, approximately 9 cm from the animal's pinna. The non-test ear was plugged with a foam insert earplug. Stimuli were

presented in ascending order of frequency (in octave steps from 0.5 kHz to 16 kHz) and intensity (in 5 dB steps). Responses (electrical activity from the recording electrode in the IC contralateral to the test ear) were amplified (20,000X) and filtered (10-3000 Hz) by a Grass P511 bioamplifier and converted to digital signals by an A/D converter on a separate DSP board in the computer. Responses were computer averaged for 100 stimulus presentations. Threshold was defined as the mid-point between the level at which there was a clear deflection in the waveform and the next lower level at which there was none.

IC-EVPs were recorded (a) prior to noise exposure in order to establish pre-exposure baselines, (b) during the conditioning exposure in order to monitor temporary TS, (c) 5 days after conditioning to document TS recovery, (d) 15 min, 24 hr, and 5 days after impulse noise exposure in order to monitor temporary TS, and (e) after 20 days recovery from high-level exposure to determine PTS. Pre-exposure and PTS measures for each animal represent the average of three tests performed on separate days. Threshold shifts were calculated by subtracting mean pre-exposure IC-EVP thresholds from post-exposure thresholds.

C. Noises and noise exposures

Animals were exposed to 0.5 kHz OBN for 5 days (6 h/day) at a level of 95 dB SPL, followed 5 days later by impulse noise simulating M16 rifle fire (Danielson et al., 1991; Henselman et al., 1994; McFadden et al., 1999). The 0.5 kHz OBN was digitally generated, low-pass filtered (TDK HAF0030 active filter set at 20 kHz), manually attenuated (Hewlett Packard 350D), amplified (NAD 2200), and delivered to a compression driver (JBL 2446J) fitted to a bi-radial exponential horn (JBL 2360H). The driver/horn assembly was suspended from the ceiling of a sound booth. Animals were housed in separate cages (approximately 27 cm X 21 cm X 22

cm) placed beneath the loudspeaker, and provided free access to food and water during noise exposure. For acoustic calibration, sound pressure levels were measured with a calibrated Type I precision sound level meter (Larson-Davis 800B) and a 1/2" condenser microphone positioned at a height corresponding to the level of the ear canal of a standing chinchilla. SPL measurements were averaged across 5 positions within each cage (geometric center and each corner). Attenuator settings and cage positions were adjusted so that the average SPL in each cage was within 1 dB of the specified SPL. Animals were rotated to different cages each day to minimize any effects of slight differences in SPL between cages.

The impulse noise was generated digitally, converted to an analog signal by a 16-bit D/A converter on a DSP board, attenuated (Hewlett-Packard 350D), amplified (NAD 2200) and routed in parallel to two compression drivers (JBL 2446) in a sound booth. Each driver was fitted with a sound delivery tube (5 cm diameter X 20 cm) with its end angled at 45° to broaden its range of resonance (Danielson et al., 1991). The ends of the sound delivery tubes were separated by 10 cm. The animal in a restraining tube was placed between the drivers with the tubes directed at the animals' ears. Each animal was exposed to 50 pairs of impulses ("salvo" exposure), with 50 ms between the two impulses of each pair and a 1000 ms interval between the onset of each pair. The total exposure time was 50 s. The impulse noise was calibrated using a 1/8" microphone (Bruel & Kjaer 4138) and voltmeter to a level corresponding to 150 dB peak SPL.

D. Cochlear analyses

After the completion of testing, chinchillas were deeply anesthetized with sodium pentobarbital (Somlethal, 400 mg/kg i.p.) and decapitated. The cochleas were quickly removed

and perfused through the oval window with a succinate dehydrogenase (SDH) staining solution consisting of 2.5 ml of 0.2 M sodium succinate, 2.5 ml of 0.2 M phosphate buffer (pH 7.6), and 5 ml of 0.1% tetranitro blue tetrazolium. Cochleas were incubated in the SDH staining solution for 45 min at 37 °C, post-fixed with 10% formalin, and stored in fixative for at least 24 hours.

Stained cochleas were dissected from the apex to the base, mounted in sections in glycerin on microscope slides, coverslipped, and examined using light microscopy (400X magnification).

The numbers of missing OHCs and IHCs were determined for successive segments of the organ of Corti. Individual cochleograms were constructed to show the percentage of hair cells missing as a function of distance from the apex of the cochlea. Percent hair cells missing was referenced to our lab standards based on normal chinchillas. Percent distance from the apex was converted to frequency using the frequency-place map of Greenwood (1990).

E. Data analysis

One male and two female chinchillas lost one or both IC recording electrodes during the experiment. Data from these ears were excluded from analysis. The final sample consisted of data from 11 ears of females and 13 ears of males. Analyses of variance (ANOVAs, SPSS) were used to assess differences between means, with IC-EVP thresholds and threshold shifts as dependent variables. Independent variables were Sex (a between-subjects factor), Frequency and Time (within-subjects factors). Significant main effects and interactions involving Sex were analyzed further using independent Student *t*-tests. Changes as a function of time were assessed using paired *t*-tests. In order to determine if sound conditioning produced significant protection from impulse noise, data obtained from animals in this experiment were compared to data obtained from a control group of 9 female and 9 male chinchillas exposed to impulse noise alone under identical conditions, described in a previous publication (McFadden et al., 1999).

Comparisons between groups were made using separate two-way Group X Sex ANOVAs for mean low-frequency PTS (average of PTS values at 0.5, 1 and 2 kHz) and high-frequency PTS (average of PTS values at 4, 8 and 16 kHz).

II. RESULTS

A. Pre-exposure thresholds

Pre-exposure thresholds are shown in Figure 1. A two-way (Sex X Frequency) mixed ANOVA showed a significant Sex X Frequency interaction, $F(5,110)=2.62$, $p = 0.028$, due to the reversal of threshold differences between low and high frequencies. Student *t*-tests at each frequency showed that the difference between females and males at 16 kHz was statistically significant, $t(22)=2.68$, $p=0.014$. Pre-exposure thresholds were similar to those of control animals in our previous study (McFadden et al., 1999).

B. Threshold shifts from conditioning noise

Threshold shifts due to sound conditioning are shown in Figure 2. After the first day of sound conditioning, thresholds were elevated by approximately 30-40 dB SPL at low frequencies, and by 0-25 dB at high frequencies. Thresholds were significantly elevated at all frequencies for females and at all frequencies except 16 kHz for males (paired *t*-tests; all p values < 0.04). Two-way (Sex X Time) mixed ANOVAs for TS at each frequency yielded significant main effects of Sex at 1, 2, 4 and 16 kHz ($F(1,22) = 5.17, 6.64, 7.02$, and 8.35 , respectively; all p values < 0.04) and main effects of Time at all frequencies except 16 kHz (all p values < 0.02). As shown in Figure 2, the main effect of Sex arose because females consistently showed more TS than males during and after the conditioning exposure.

Paired *t*-tests were used to follow-up on the significant main effects of Time. These analyses indicated that TS decreased significantly at 0.5 kHz between Day 1 and Day 5 of conditioning for both males, $t(12)=2.84$, $p = 0.015$, and females, $t(10)=2.97$, $p=0.014$. The decreases in TS that occurred between the last day of conditioning and the end of the 5-day recovery period were significant at all frequencies except 16 kHz (all p values < 0.04). After 5 days of recovery from conditioning, thresholds were within approximately 5 dB of pre-exposure values at all frequencies (Fig. 2, bottom panel).

C. Threshold shifts after impulse noise

Mean TS values at 15 min, 24 h and 5 days after impulse noise exposure are shown in Figure 3. Two-way (Sex X Time) mixed ANOVAs for TS at each frequency showed significant Sex X Time interactions at 2 and 4 kHz ($F(3,66) = 5.36$ and 3.14 , respectively, p values < 0.04), and significant main effects of Time at all frequencies (all p values < 0.001). As shown in Figure 3, the interactions occurred because females showed more recovery at 2 and 4 kHz over time than males.

Mean PTS is shown in Figure 4. Paired *t*-tests indicated that impulse noise produced significant PTS at all frequencies for both males and females (all p values ≤ 0.01). A two-way (Sex X Frequency) ANOVA showed a significant Sex X Frequency interaction, $F(5,110)=4.20$, $p = 0.002$. The interaction occurred because males developed more PTS than females at low frequencies, but less PTS than females at 16 kHz.

To summarize the IC-EVP test results, females had a significantly lower mean threshold at 16 kHz than males prior to exposure (Fig. 1). During conditioning, females consistently showed greater TS than males, but thresholds of both sexes were essentially normal within 5 days after

conditioning (Fig. 2). Subsequent exposure to M16 rifle fire produced more TS at low-frequencies for males, and more TS at high frequencies for females (Fig. 3). When PTS was assessed 20 days after exposure to M16 rifle fire, females showed approximately 5-10 less PTS than males at 0.5, 1 and 2 kHz, but approximately 15 dB more PTS than males at 16 kHz.

D. Noise-induced hair cell losses

Mean cochleograms are shown in Figure 5. Hair cell lesions were slightly smaller in cochleas from females. Average OHC losses in the apical half of the cochlea were approximately 30% for females and 40% for males. Average OHC losses in the basal half were approximately 50% for females and 60% for males. The most striking difference was seen in the 2 kHz region of the cochlea, where males showed approximately 30% more OHC loss than females. Mean IHC losses were not remarkably different between females and males. Averaged across the entire cochlea, females had 11% IHC loss and 38% OHC loss, whereas males had 14% IHC loss and 47% OHC loss.

E. Protection from M16 rifle fire

One purpose of exposing animals to the 5-day conditioning regimen was to provide protection from subsequent exposure to M16 rifle fire. A perspective on the success of this approach is provided by Figure 6, which shows differences in PTS between conditioned animals in the present study and control animals exposed only to the impulse noise (McFadden et al., 1999). Sound conditioning provided up to 18 dB protection for females and up to 10 dB protection for males at individual frequencies. Collapsed across sex, the protective effect was 5 to 12 dB across frequencies, with greater protection at low frequencies than at high frequencies.

Figure 7 compares average low-frequency PTS and high-frequency PTS for conditioned and control animals. The pattern of PTS was similar for conditioned animals and controls. For both groups, females showed less low-frequency PTS but more high-frequency PTS than males. Conditioned females and males both showed less PTS than their same-sex controls. Two-way (Group X Sex) ANOVAs showed significant main effects of Group, $F(1,51)=6.70$, $p=0.012$, and Sex, $F(1,51)=4.70$, $p=0.035$, for low-frequency PTS. Thus, conditioned animals showed significantly less low-frequency PTS than controls, and females showed significantly less low-frequency PTS than males. Despite the consistent trends for high-frequency PTS, neither differences between males and females nor differences between conditioned animals and controls were statistically significant.

Hair cell lesions were smaller in the conditioned animals compared to controls. Conditioned males had approximately 20% less IHC loss and 24% less OHC loss than control males. Conditioned females had approximately 5% less IHC loss and 18% less OHC than control females. Collapsed across sex, sound conditioning resulted in approximately 12% less IHC loss and 21% less OHC loss compared to controls.

III. DISCUSSION

A. Protective effects of sound conditioning

The results show that sound conditioning provides protection from PTS and hair cell loss caused by impulse noise exposure. Chinchillas conditioned with a 0.5 kHz OBN at 95 dB SPL for 6 h/day for 5 days developed approximately 5-12 dB less PTS across frequencies, and 13-21% less hair cell loss than chinchillas exposed to the impulse noise alone. As shown in Figure 7, protection was apparent at high frequencies (approximately 6 dB overall) as well as low

frequencies (approximately 10 dB overall). However, only the protection at low frequencies reached statistical significance.

The present results confirm and extend the results of an earlier study by Henselman et al. (1994) by showing that a 5-day conditioning regimen can protect the ear from impulse noise. A comparison between the two studies also shows a "dose effect" related to the duration of the conditioning exposure. In the Henselman experiment, a 10-day conditioning regimen produced approximately 7-23 dB of protection across frequencies, with the greatest protection (20-23 dB) at 2 and 4 kHz. Averaged across frequencies, the protective effect of a 10-day conditioning regimen was approximately 16 dB SPL. The 5-day conditioning regimen used in the present study also provided significant protection from PTS. However, the magnitude of protection was approximately 6-10 dB less than that provided by the longer conditioning exposure. Differences related to the duration of the conditioning exposure are also apparent in hair cell losses. The pattern of hair cell loss we observed in the present study was similar to the pattern seen by Henselman et al. (1994). However, the magnitude of hair cell protection was considerably greater for the 10-day conditioning regimen. Animals conditioned for 5 days (present study) had approximately 40% OHC loss across the entire cochlea, whereas animals conditioned for 10 days (Henselman et al., 1994) had less than 20% OHC loss. The "dose effect" of conditioning is interesting because it indicates that the magnitude of protection can be increased by extending the "training" period. The mechanisms by which sound conditioning increases resistance to NIHL are not known, but they are clearly not "all or none" phenomena. Previous studies have indicated dose effects of sound conditioning related to exposure duration (Canlon and Fransson, 1998; Subramaniam et al., 1993) and rest period between conditioning and high level exposure (McFadden et al., 1997; Subramaniam et al., 1992).

It is interesting that sound conditioning can protect the cochlea from impulse noise, which can damage the cochlea by causing direct mechanical failure as well as through metabolic disruption. Whether sound conditioning actually increases resistance to mechanical damage, or whether it only attenuates damage brought about by metabolic changes is not clear. However, there is some evidence that the structural components of the ear may be altered by sound conditioning in a way that could afford protection from mechanical damage (Hu and Henderson, 1997; Pack et al., 1999).

B. Differences between female and male chinchillas

The results show significant differences between female and male chinchillas in their response to noise. During sound conditioning, females consistently showed greater TS than males. After impulse noise exposure, females developed approximately 10 dB less PTS at low frequencies, but approximately 5 dB more PTS at high frequencies than males. The difference in low-frequency PTS between females and males was statistically significant. The pattern of PTS in sound conditioned animals was very similar to that reported previously for animals exposed to impulse noise alone (McFadden et al., 1999). In both cases, females had less TS at low frequencies and greater TS at high frequencies than males. Sound conditioning provided significant protection for both females and males, with little difference in the magnitude of protection.

The reasons for the sex differences cannot be determined from this study. However, since the differences were observed in chinchillas, extraneous factors such as differences in noise exposure history, recreational activities, and dietary factors can be ruled out. One possibility is that sex/gender differences in susceptibility to NIHL arise from differences in the acoustical

properties of the outer and middle ears. A second possibility is that sex/gender differences arise from basic physiological differences between female and male cochleas. Interestingly, Mills et al. (1999) recently reported that male rats are more susceptible to kanamycin ototoxicity than female rats, a difference that clearly cannot be attributed to differences in acoustical properties of the ear. McFadden et al. (1998) observed changes in otoacoustic emissions in a human male treated with estrogen, indicating that sex hormones can influence outer hair cell function. Because the outer hair cell system is a likely candidate as a site for conditioning-induced changes (Canlon, 1996; Canlon and Fransson, 1995; Hu and Henderson, 1997), observations such as this may be important for understanding the sex differences we have observed in chinchillas. Future studies using the chinchilla may help determine the relative importance of anatomical and physiological factors in sex/gender differences in sound conditioning and susceptibility to NIHL.

ACKNOWLEDGMENTS

This research was supported by Grant DAMD17-96-1-6330 to SLM. The authors are grateful to Marty Howard for her assistance in data collection and noise exposures.

Figure Legends

Figure 1. Mean thresholds of female (solid circles) and male chinchillas (open triangles) before noise exposure. Bars in this and subsequent figures represent standard errors of the means (SEM).

Figure 2. Mean threshold shifts (\pm SEM) during and after sound conditioning. Top panel: threshold shifts after the first day (6 h) of sound conditioning. Middle panel: threshold shifts after the last day of conditioning. Bottom panel: threshold shifts 5 days after the last conditioning exposure and before exposure to impulse noise.

Figure 3. Mean threshold shifts (\pm SEM) at three times after impulse noise exposure. Top panel: threshold shifts 15 min after impulse noise exposure. Middle panel: threshold shifts one day after impulse noise exposure. Bottom panel: threshold shifts 5 days after impulse noise exposure.

Figure 4. Permanent threshold shifts (PTS) measured 20 days after impulse noise exposure for females (solid circles) and males (open triangles). Bars show SEMs.

Figure 5. Cochleograms showing hair cell loss after impulse noise exposure. Left panel: outer hair cell (OHC) losses for males (solid lines) and females (dashed lines). Right panel: inner hair cell (IHC) losses for males (solid lines) and females (dashed lines).

Figure 6. Protection from permanent noise-induced hearing loss afforded by sound conditioning in females (solid circles) and males (open triangles). Protection is the difference in permanent threshold shift (PTS) between control animals exposed to the impulse noise alone and animals conditioned with 0.5 kHz octave band noise at 95 dB SPL for 5 days (6 h/day).

Figure 7. Mean permanent threshold shifts (PTS) at low frequencies (average of 0.5, 1 and 2 kHz) and high frequencies (average of 4, 8 and 16 kHz) for conditioned females and males and their same-sex controls. Bars show SEMs.




Figure 1




Figure 2




Figure 3




Figure 4







Figure 5




Figure 6




Figure 7